




Winter Outlook 2016 – 2017

National Weather Service Pendleton Oregon
And NOAA Climate Prediction Center (CPC)




El Nino-Southern Oscillation (ENSO)

- ENSO-neutral conditions are present
- Equatorial sea surface temperatures (SST) are near or below average in the east-central & east Pacific Ocean
- ENSO-neutral conditions are slightly favored (55-60%) during the upcoming NH fall/winter 2016-17

Average SST Anomalies
21 AUG 2016 – 17 SEP 2016





EQ. Upper-Ocean Heat Anoms. (deg C) for 180–100W




ENSO- Outlook (CFSV2)

The CFS.v2 ensemble mean (black dashed line) generally favors borderline ENSO-neutral/ La Niña during the Northern Hemisphere fall and winter 2016-17.



Official CPC Temperature Outlook Dec-Jan-Feb

- The Climate Prediction Center, or CPC is calling of equal chances of above, below or near average temperatures through the three month period (Dec-Jan-Feb) over much of the Pacific Northwest. Southeastern Oregon, southern Idaho and the remainder of the Southwest does have higher probabilities for above average temperatures during this time period. Please remember that weekly, and daily temperatures will still fluctuate from above, to below average at times.



Local 3 Month Temperature Outlook: Yakima, WA

Three-Month Temperature Outlook (Issued: September 2016)

[National Outlook](#)

[Local Outlook](#)

[Background Information](#)

[Questions and Feedback](#)

YAKIMA MUNICIPAL AP, WA

YAKIMA County, Coop ID: 459465


Elevation: 1064 ft.

Latitude: 46° 34' N Longitude: 120° 32' W

[NOAA Online Weather Data](#)

[Help](#)

Three Category Temperature Outlook: Dec-Jan-Feb 2016_17



There is a 33.0% chance for the average temperature during this 3-month period to be higher than 33°.

There is a 34.0% chance for the average temperature during this 3-month period to be between 30.3° and 33°.


There is a 33.0% chance for the average temperature during this 3-month period to be lower than 30.3°.

*Outlooks are calculated using the 1981-2010 climatological reference period.

[Table Text Only](#)


Local 3 Month Temperature Outlook: Kennewick, WA

Three-Month Temperature Outlook (Issued: September 2016)




Local 3 Month Temperature Outlook: Bend, OR

Three-Month Temperature Outlook (Issued: September 2016)



Official CPC Precipitation Outlook Dec-Jan-Feb




The Climate Prediction Center (CPC) is calling for equal chances for above, below or near average precipitation totals during the three month period across Central Oregon, north to Central and Western Washington. The probabilities are tilted slightly in favor of higher than average precipitation totals over Northeast Oregon, and much of Eastern Washington. The southern tier of the US, from Arizona to Florida has greater odds of below average precipitation totals during this time. Above average precipitation totals are favored in the Great Lakes, where abundant lake effect snow may fall this year.






How are these CPC Outlooks Produced?

- Section 1: CFSV2 forecasts for November, December, January, February and the combined three month Dec-Feb forecast.
- Forecasts include 700mb height anomalies (upper left)...forecast surface temperature anomalies (upper right) and precipitation anomalies (bottom)




CFSv2 November



CFSv2 December




CFSv2 January




CFSv2 February


CFSv2 monthly z700 anomalies (m; shaded) and total (dm; contours)




CFSv2 monthly T2m anomalies (K)



CFSv2 monthly Prec anomalies (mm/day)




CFSv2 Dec-Jan-Feb




Box & Whiskers Temperature: Columbia Basin

DJF Temperature Distribution for Climate Div. #074



Box & Whiskers Precipitation: Columbia Basin

DJF Precipitation Distribution for Climate Div. #074




Next: Analog Year Plots



- 500mb Height Anomalies
- Surface Temperature Anomalies
- Precipitation Anomalies
- Average of past years that also had a transition from a strong El Nino to very weak La Nina or neutral state in the tropical Pacific (8 years total)

December

NCEP/NCAR Reanalysis
500mb Geopotential Height (m) Composite Anomaly 1981-2010 climo




NCEP/NCAR Reanalysis
Surface air (C) Composite Anomaly 1981-2010 climo




January


NCEP/NCAR Reanalysis
500mb Geopotential Height (m) Composite Anomaly 1981–2010 climo



NCEP/NCAR Reanalysis
Surface air (C) Composite Anomaly 1981–2010 climo




NCEP/NCAR Reanalysis
Surface Precipitation Rate (mm/day) Composite Anomaly 1981–2010 climo




February


NCEP/NCAR Reanalysis
500mb Geopotential Height (m) Composite Anomaly 1981–2010 climo



NCEP/NCAR Reanalysis
Surface air (C) Composite Anomaly 1981–2010 climo




NCEP/NCAR Reanalysis
Surface Precipitation Rate (mm/day) Composite Anomaly 1981–2010 climo




Dec-Jan-Feb


NCEP/NCAR Reanalysis
500mb Geopotential Height (m) Composite Anomaly 1981-2010 climo



NCEP/NCAR Reanalysis
Surface air (C) Composite Anomaly 1981-2010 climo



NCEP/NCAR Reanalysis
Surface Precipitation Rate (mm/day) Composite Anomaly 1981-2010 climo



Summary

- There can be other factors not mentioned in this presentation that affect our winter weather in the Pacific Northwest (PDO, PNA, AO and even NAO indices). Often times, on a more of a weekly or bi-weekly time scale.
- The weather patterns throughout the winter months will undoubtedly fluctuate and change. Bringing periods of above, below and near normal temperatures and precipitation.
- The official Climate Prediction Center(CPC) outlooks for December – February are calling for equal chances of above, below or near average temperatures for much of the area. Meanwhile precipitation outlooks for the same period have the odds slightly tilted toward above average precipitation totals, especially over Northeast Oregon, and much of Eastern Washington.